



This is a repository copy of *Understanding the factors that affect retention within the mental health nursing workforce: a systematic review and thematic synthesis*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/175874/>

Version: Published Version

---

**Article:**

Adams, R., Ryan, T. and Wood, E. (2021) Understanding the factors that affect retention within the mental health nursing workforce: a systematic review and thematic synthesis. *International Journal of Mental Health Nursing*, 30 (6). pp. 1476-1497. ISSN 1445-8330

<https://doi.org/10.1111/inm.12904>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



REVIEW ARTICLE

# Understanding the factors that affect retention within the mental health nursing workforce: a systematic review and thematic synthesis

Rosie Adams, Tony Ryan and Emily Wood

Health Sciences School, The Division of Nursing and Midwifery, The University of Sheffield, Sheffield, UK

**ABSTRACT:** *There are over 41 000 vacant nursing posts across the United Kingdom's National Health Service (NHS), with more people leaving the profession than joining it. Despite mental health being acknowledged as a priority area, some of the most significant staff shortages are occurring within mental health services. Urgent action is needed to retain the mental health nurses (MHNs) currently in post to ensure the profession is fit for purpose and aid future recruitment efforts. This review set out to identify the individual factors that affect the retention of MHNs. A systematic search of six databases was conducted (CINAHL, PsychINFO, MEDLINE, Web of Science (Core collection), EMBASE and the British Nursing Index). Studies were systematically screened for inclusion based on predetermined eligibility criteria. The studies were quality assessed using the Mixed Methods Appraisal Tool (MMAT). Findings were synthesized using Thematic Synthesis. A total of 23 studies consisting of a range of study designs were included in the review. Four key themes emerged from the synthesis: Individual characteristics, Working within mental health services, Training and skills and Work environment. The findings from this review suggest that MHNs encounter some factors unique to working in mental health services, which suggests that retention strategies should be specific to each nursing speciality. Beyond nursing speciality, the factors identified vary between clinical settings in mental health due to the differences in work environments and services they provide. Future studies should now set out to explore what factors exist in which clinical settings to inform better tailored retention strategies to generate better outcomes.*

**KEY WORDS:** *intent to leave, intent to stay, mental health nurse, retention, RMN.*

## BACKGROUND

Nursing shortages represent a global concern, with the World Health Organization (2013) estimating a 12.9

million deficit in skilled health professionals made up of physicians, nurses and midwives by 2035. The most significant shortages are currently occurring in Africa and South-East Asia (World Health Organization 2020), whilst future nursing workforces in the UK, United States, Australia and Portugal are vulnerable to significant projected shortages from an ageing workforce and limited workforce planning policies (Buchan *et al.* 2015). Nurses make up 50% of the global health-care workforce (World Health Organization 2020) and account for over a quarter of the UK's NHS staff (Beech *et al.* 2019). With increasing demand, insufficient resources and deteriorating job satisfaction, the NHS nursing workforce is overstretched and depleting rapidly, now accounting for 40% of staff shortages in

**Correspondence:** Rosie Adams, The University of Sheffield, Health Sciences School, The division of Nursing and Midwifery, Barber House Annexe, 3a Clarkehouse road, Sheffield, S10 2LA, UK. Email: raadams1@sheffield.ac.uk

**Authorship statement:** Rosie Adams drafted the paper, conducted the search, data extraction, quality assessment, synthesis and discussion of the findings. Emily wood and Tony Ryan assisted the search, and contributed to the synthesis and discussion of the findings. All authors have read and approved the final manuscript.

**Declaration of Conflict of interest:** None.

Rosie Adams, MSc, RMN.

Tony Ryan, PhD.

Emily Wood, PhD, RN.

Accepted June 15 2021.

© 2021 The Authors. *International Journal of Mental Health Nursing* published by John Wiley & Sons Australia, Ltd.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

hospital, community and mental health services (Beech *et al.* 2019). There are currently over 41,000 vacant nursing posts within the NHS (Buchan *et al.* 2019), with more people currently leaving the profession than joining it (Nursing & Midwifery Council 2017). Despite efforts to increase the number of nurses being trained, there was a decline in the numbers of nursing applications for nurse training in England in 2017 and 2018 (Buchan *et al.* 2019).

Mental health is a priority area identified within the 'NHS long-term plan', 'interim NHS people plan' and 'five-year forward view', yet the most significant staff shortages are occurring within the NHS mental health nursing workforce. Years of underinvestment and funding cuts to mental health services, nurses pay and training bursaries have given rise to these staff shortages (Unison 2017a), evidenced by a drastic 12% fall in MHNs between 2010 and 2017 in the NHS (Care Quality Commission 2017). Even with increased efforts to recruit and retain MHNs, their numbers rose by <0.5% in 2018 (Buchan *et al.* 2019). Due to poor retention and high turnover, a National Retention Programme (NRP) was launched in 2017 by the NHS. Under this scheme, clinical mental health staff turnover has decreased from 14.3% to 13.4%, but there is still a long way to go to resolve this current crisis (NHS Improvement 2019a). Poor retention reduces the ability of MHNs to meet service user's needs leading to dangerous consequences for service user care and safety (Unison 2017a) and is likely to deter people from wanting to join the profession.

Developing strategies focused on encouraging recruitment may be a futile occupation if retention is not addressed. Whilst nurse turnover can be beneficial by bringing in new perspectives, ideas and experience, high turnover rates often come with high economic and non-economic costs (Dewanto & Wardhani 2018). Instability of nursing staff leads to unstable nursing care, which increases the risk of care and treatment errors that can have grave effects of patient care and safety (O'Brien-Pallas *et al.* 2010), whereas relying on bank and agency staff, continuous recruitment and orientation and training processes constitutes a large economic burden (Jones & Gates 2007). In 2018, the NHS had reportedly spent £353 million on agency and bank staff to meet safe staffing requirements (Buchan *et al.* 2019). Urgent action is required to reverse this trend and reduce the use of bank and agency nurses (Royal College of Nursing 2015), before patient safety is put at greater risk

(Unison 2017b). More immediate emphasis should be placed on retaining the staff already in post. By doing this, we can help to ensure the current profession is fit for purpose, and one that others aspire to join to aid retention and boost recruitment (Unison 2017b).

Job dissatisfaction appears to be strongly associated with poor retention (Happell 2008; Health Education England 2014; Ward 2011). Job satisfaction for MHNs is reportedly the lowest it has ever been, with many nurses emotionally and physically exhausted, leading them to consider leaving their profession (Unison 2017b). This results in a negative cycle where the consequences of poor job satisfaction and turnover, for example increased workload, pressure and working with temporary or inadequate numbers of MHNs, lead to even more turnover (Jones & Gates 2007). Recent events have seen us delve into the depths of a global pandemic which has now placed even more pressure on the mental health workforce, due to increasing demand and even more staff shortages as a result of shielding, sickness and self-isolation guidance (British Medical Association 2020). The additional pressures of the pandemic are likely to result in an exacerbation of the already significant international nurse shortages (Turale & Nantsupawat 2021).

Much of the literature surrounding nurse turnover and retention is for adult nursing (Chan *et al.* 2013; Halter *et al.* 2017; Moseley *et al.* 2008). Whilst there is some research focused on MHN retention, there has not yet been a systematic review consolidating the evidence on factors that affect retention within this workforce. With the concerning consequences of poor retention on patient care and safety, it is imperative to get a better understanding of the specific factors causing MHNs to leave or contemplating leaving; so that meaningful and targeted retention strategies can be put in place to resolve this current crisis. The aim of this review is to identify what factors affect the retention of MHNs, in order to inform retention strategies and identify areas for future research.

## METHODS

### Protocol and registration

This systematic review and thematic synthesis adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines (Moher *et al.* 2009).

## Database searches

A systematic search of six databases was conducted from their year of inception to November 2019: CINAHL, PsychINFO, MEDLINE, Web of Science (Core collection), EMBASE and the British Nursing Index (BNI). Final search strategies were developed using the following key terms: ("Mental health nurs\*" OR "psychiatric nurs\*" OR RMN\* OR "mental nurs\*" OR RMHN) AND (Retention OR retain\* OR remain\* OR leav\* OR attrition OR "intent\* to leave" OR "intent\* to stay" OR turnover OR "sustainable workforce" OR loyalty OR resign\* OR shortage). No date limits were applied to ensure a sufficient number of results.

## Grey literature and other resources

Although the BNI includes dissertations and theses, it mainly covers papers published within the UK. Therefore, to supplement the electronic searches and minimize publication bias, grey literature databases were searched for any relevant unpublished studies including dissertations and theses. Reference lists from relevant studies and reviews were also screened for eligible studies.

## Contacting authors

Primary authors were contacted to try and gain the full-text versions of unpublished articles that could not be located elsewhere.

## Inclusion criteria

The following inclusion criteria were used to identify eligible studies:

### Population

- Studies focussing on registered nurses who specialize in mental health or work in mental health settings of any age, experience or staff grade.

### Setting

- Studies focussed on registered nurses working in any mental health setting.

### Outcomes

- Studies reporting any findings on factors that affect the retention of registered nurses working within mental health settings.

- Studies reporting findings on interventions or strategies that aim to affect retention.
- Studies reporting findings on the predictive factors of retention.

### Study design

- Any empirical studies including all study designs.
- Studies were not excluded based on quality due to the paucity of research on this topic.
- Studies not written in the English language at full text were excluded.

## Study selection

The search results generated from the databases were exported to Mendeley, and duplications were removed. Titles and abstracts were screened simultaneously to identify relevant papers. Relevant papers were then screened at full text to identify eligible studies for review based on the predetermined eligibility criteria.

## Data extraction

A simple standardized data extraction form was created based on the information required to answer the review question using Excel. The extracted data were tabulated and assessed for heterogeneity.

## Quality assessment

As the included studies comprised a range of study designs and methodologies, the MMAT tool version 2018 and user guide were used to assess the quality of the studies (Hong *et al.* 2018). The included studies were categorized and rated using the appropriate quality criteria. These ratings were then used to form a percentage score. Each of the quality criteria within each category will account for 20% of an overall score, so if a study met all the criteria they would score 100%. Scores were accompanied by a summary of the quality by comparing and contrasting the studies within and across categories. Mixed methods studies were rated using the quality criteria for mixed methods studies plus the appropriate quality criteria for each individual study component.

## Data synthesis

Meta-analysis was not possible due to the level of heterogeneity within the included studies. Thus, a

thematic synthesis was conducted to identify and generate descriptive themes across all included studies. Findings from quantitative studies were converted into qualitative form by a process often described as 'qualitizing' or 'data transformation' (Dixon-Woods *et al.* 2005; and Pluye & Hong 2014). This was done by transforming the statistics derived from survey/questionnaire responses and secondary data sets into textual data, in order to create a narrative of the results that can be synthesized alongside the qualitative findings (Heyvaert *et al.* 2016). The data transformation process can be seen in Appendix A in the supplementary materials. The qualitized data along with the qualitative findings were then imported to Quirkos software for analysis. Thomas and Harden's (2008) stages of thematic synthesis were followed: the process began with line-by-line coding of the findings of each study; descriptive themes were then developed through organizing the codes into themes and associating themes with similar concepts; the final themes and subthemes were accompanied by a narrative analysis and then contextualized within the existing literature in the Discussion section.

## RESULTS

### Search results and study selection

The search of the six databases retrieved 4428 potentially relevant papers. No additional papers were identified through searching grey literature or checking the reference lists of relevant papers. After duplicates were merged, the citations and abstracts were screened simultaneously for 2024 papers. This left 45 papers eligible for screening at full text. Primary authors for two of the studies were contacted to access the full-text version of their papers for screening which were unable to be located otherwise (Hanohano & Dee 2017; Walrath 2011). Walrath provided the full-text version for screening. Hanohano and Dee's (2017) paper was excluded at full text as it was unavailable. A total of 22 papers were excluded at full text with reasons. This left 23 papers eligible for inclusion; see Figure 1 for an overview of the study selection process.

### Description of the included studies

The detailed characteristics for the twenty-three included studies can be seen in Table 1. Of the studies, twenty were published in journals and peer-

reviewed nursing magazines, and three were unpublished PhD theses. The studies span almost three decades from 1991 to 2019 including 18 918 MHNs. The studies were conducted in eight different countries, nine in the United States, six in the United Kingdom, three in Jordan, and one in Canada, China, Japan, Ghana and Israel. Although a lot of the same factors affecting retention are identified in studies conducted in different countries, it is important to note that there are differences in the nurse training programmes and healthcare systems in different countries that generated different factors for nurses working in different countries.

The clinical settings included in the studies comprised of NHS, private, state-run and Ministry of Health settings. Over half of the studies were conducted in hospitals including wide a range of inpatient settings, and six studies included a range of community-based settings which can be seen in Table 1. Six studies did not specify the individual clinical settings (Hamaideh 2011; Murrels & Robinson 1999; Robinson *et al.* 2005; Robinson & Tingle 2003; Sherring & Knight 2009; Walrath 2011).

The studies included a range of study designs. Two studies were evaluating retention strategies. Rollins (2014) evaluated whether an increase in the length and content of an existing training programme undertaken in MHNs' orientation period impacted retention. The new 'healthy work environment' (HWE) programme was extended to eight weeks and included familiarization to local polices and documentation, skill competencies and seclusion and restraint training. Pelletier *et al.* (2019) introduced a nurse residency programme for newly graduated nurses. The programme incorporated training on a vast number of clinical theories and practice, as well as topics that influence retention derived from the literature, and provided a social support system.

The aims of the studies varied with the majority of the studies ( $n = 18$ ) directly aiming to explore factors that affect retention, although the factors explored differed among the studies. The rest of the studies ( $n = 5$ ) had different aims with different foci that can be seen in Table 1. These studies were included as they still generated some important findings on factors that affect retention. Due to the differences in study design and aims, the outcome measures also varied significantly. All the qualitative studies used individual interviews and the quantitative studies used a range of validated and unvalidated questionnaires to explore different factors that affect retention.



**FIG. 1** Study selection flow diagram.

## Quality assessment

The included studies were categorized into 4 categories of study designs: 'qualitative' ( $n = 4$ ) (Alexander *et al.* 2015; Karlowicz & Ternus 2009; Musto & Schreiber 2012; Stacey *et al.* 2011), 'quantitative non-randomized' ( $n = 2$ ) (Pelletier *et al.* 2019; Rollins 2014), 'quantitative descriptive' ( $n = 15$ ) (Alsaraireh *et al.* 2014; Bamber 1991; Baum & Kagan 2015; Hamaideh 2011, 2014; Ito *et al.* 2001; Jiang *et al.* 2019; Kagwe *et al.* 2019; Murrels & Robinson 1999; Robinson & Tingle 2003; Robinson *et al.* 2005; Sherring & Knight 2009; Walrath

2011; Yanchus *et al.* 2015, 2017) and 'mixed methods' ( $n = 2$ ) (Agyapong *et al.* 2015; Gunn 2015). The details of the individual assessments for each individual study can be seen in Appendix B in the supplementary materials. All of the qualitative studies met 100% of the quality criteria indicating they are high-quality studies. One of the quantitative non-randomized studies met 80% of the quality criteria but did not report on possible confounders (Rollins 2014), whereas Pelletier *et al.*'s (2019) quantitative non-randomized study only met 60% of the quality criteria. This study had a response rate of 33% due to attrition over time. Five of

**TABLE 1** *Study characteristics*

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
Alexander et al (2015)	To explore how nurses came to choose and remain in mental health nursing careers	Qualitative Descriptive Phenomenology	United States	Acute inpatient settings	MHNs who have worked in acute inpatient settings for 5 years or longer. <i>n</i> = 8	1 hour face-to-face, semi-structured interviews	Stereotypes and career pride Positive team dynamics Remaining hopeful Leadership Moral distress
Musto and Schreiber (2012)	To develop a theory of the process used by mental health nurses when they experience moral distress	Qualitative Grounded theory	Canada	Inpatient units and community settings for adolescents with mental health issues.	RNs who work in the included clinical settings. <i>n</i> = 12	60–120 min semi-structured interviews.	
Stacey et al (2011)	To explore how values influence the experience of nursing practice for nurses working in inpatient setting	Qualitative Narrative	United Kingdom	Adult inpatient settings	MHNs with between 6 months' and 3 years' experience in adult inpatient settings. <i>n</i> = 12	One-to-one interviews	Conflict of values in practice
Karlowicz and Ternus (2009)	To gain understanding of and describe psychiatric nurses' experiences within their first year of employment, and identify the factors that influenced their decision to remain in or leave the organization	Qualitative Case analysis	United States	Inpatient psychiatric services	MHNs employed by the organization and those who left the organization within the first year of employment. <i>n</i> = 14	Telephone interviews	Role and identity Orientation Education and training Organizational expectations Team dynamics Leadership Salary
Rollins (2014)	To determine how recent changes to the length and content of the psychiatric nurse orientation training impacted psychiatric nurse retention rates	Quantitative non-randomized Pre- and post-intervention design PhD Thesis	United States	Inpatient Outpatient Residential Partial Day	All RNs working within the 4 clinical settings. <i>n</i> = 88	Human Resources data set for retention rates pre and post-intervention.	Healthy Work Environment Orientation training
Pelletier et al (2019)	To examine the effectiveness of a new nurse residency programme for retaining new graduate nurses in a mental health setting	Quantitative non-randomized Comparative study design	United States	Inpatient units	New graduate nurses hired into the new nurse graduate programme <i>n</i> = 34	Human Resources data set to measure turnover rates Job/Work environment Nursing Satisfaction Survey Eisenberger Social Support Scale OCB scale CIV ACS	Nurse Residency Programme

(Continued)

**TABLE 1** (Continued)

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
Alsaraireh et al. (2014)	"To fill a gap in the understanding of psychiatric nursing in Jordan and provide information for future human resource planning"	Quantitative descriptive Cross-sectional	Jordan	A governmental hospital for mental health consisting of 1 male acute and long-term unit, and 1 female acute and long-term unit.	Nurses working in the hospital. <i>n</i> = 154	POF WBI CSE Occupational commitment tool MSQ-Short form to measure job satisfaction. WSC to measure turnover intention.	Job satisfaction Gender Marital status Nurse training experience Salary Different inpatient settings Age
Jiang et al (2019)	"To investigate the intention to leave among psychiatric nurses in China, and to identify associated factors"	Quantitative descriptive Cross-sectional	China	Tertiary psychiatric hospitals	MHNs working in tertiary psychiatric hospitals. <i>n</i> = 7933	MSQ-Short form (Weiss <i>et al</i> , 1967) to measure job satisfaction. Researcher-developed questionnaire to measure individual characteristics, job-related variables, and intention to leave.	Gender Age Marital status Level of education Amount of working hours Time in job Patient-initiated violence Health Staff grade Salary Patient respect Self-recognition Physician nurse coordination and trust Job satisfaction
Walrath (2011)	"To examine the correlation, if any, between nurse leaders' communication effectiveness and nurses' perceived job performance and satisfaction"	Quantitative descriptive Quantitative correlational design PhD Thesis	United States	State government treatment centre.	MHNs (including direct care and supervisory nurses) working at the psychiatric treatment centre <i>n</i> = 21	SLCI to measure supervisor leadership and employee performance. JSS (to measure communication and supervision subscales) Research developed questions regarding individual characteristics and intent to stay employed.	Supervisor leadership Supervisor communication Supervisor communication
Ito et al (2001)	To examine psychiatric nurses' intention to leave their job in relation to their job	Quantitative descriptive Cross-sectional	Japan	Psychiatric hospitals	All licenced nurses working in the included psychiatric hospitals. <i>n</i> = 1494	NIOSH job stress questionnaire to measure job satisfaction, perceived risk of	Other available opportunities Age Time in current job

(Continued)



**TABLE 1** (Continued)

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
	satisfaction, perceived risk of assault, and supervisory support					assault and supervisory support.	Previous job changes Supervisory support Perceived risk of assault Job satisfaction
Murrells and Robinson (1999)	To document career plans and histories of mental health nurses postqualification, and identify and explore factors which may impact directions followed inside and outside the NHS	Quantitative descriptive Longitudinal	United Kingdom	NHS and non-NHS settings	Last groups of students completing traditional routes to the RMN qualification prior to the new diploma (project 200) courses <i>n</i> = 447	Researcher-developed questionnaire.	Age Course preparation
Bamber (1991)	To Provide a detailed analysis of reasons for leaving among psychiatric nurses	Quantitative descriptive Two-year prospective study using survey design	United Kingdom	NHS psychiatric hospital	Staff nurses working within the psychiatric hospital <i>n</i> = 82	Researcher-developed questionnaire to measure job satisfaction. BI (Berkley Planning Associates) to measure stress. Framingham Type A Personality questionnaire to measure personality type.	Job satisfaction Perceived quality of decisions made by those in managerial positions. In-service training Physical working conditions Burnout Personality type Clinical experience Age Level of education
Robinson and Tingle (2003)	To ascertain diversity of the workforce, describe career plans and pathways postqualification, investigate experiences relevant to careers, and identify relationships between career plans, plans followed and profile and experience	Quantitative descriptive Longitudinal	United Kingdom	Various NHS psychiatric settings	Mental health diplomates <i>n</i> = 678	Researcher-developed questionnaires	Working conditions Providing care Working relationships Reflection on practice CPD opportunities
Sherring and Knight (2009)	To describe burnout and develop an understanding of the variables involved in burnout for mental health	Quantitative descriptive Cross-sectional	United Kingdom	City NHS Trust.	MHNs working at the included Trust. <i>n</i> = 172	MBI	Emotional exhaustion Depersonalization

(Continued)

**TABLE 1** (Continued)

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
Yanchus et al (2017)	nurses working in a city Trust To examine predictors of turnover intention, or employee's cognitive withdrawal from their job among direct care mental health professionals	Quantitative descriptive Cross-sectional	United States	VHA	Veterans Affairs employees including RNs, Social workers, psychologists, and psychiatrists. RNs <i>n</i> = 2432	Veterans Affairs All Employee Survey 2015 to measure: civility, emotional exhaustion, supervisory support, job satisfaction, workgroup and supervisory psychological safety, turnover intention and turnover plans.	Supervisory support Job satisfaction Emotional exhaustion Civility
Robinson et al (2005)	To ascertain diversity of the workforce, describe career plans and pathways postqualification, investigate experiences relevant to careers, and identify relationships between career plans, plans followed and profile and experience	Quantitative descriptive Longitudinal	United Kingdom	Various NHS psychiatric settings	Mental health diplomates. <i>n</i> = 678	Researcher-developed questionnaire.	Consolidation of skills Job security Time in job Working relationships/good team Staff grade Salary Career development CPD opportunities Staffing levels and ratios Clinical supervision Preceptorship Work to life balance Ethnicity Leadership support Paperwork Age Moral distress
Hamaideh (2014)	To describe the levels and predictors of moral distress of mental health nurses, and relationships of moral distress with nurses' intention to leave, burnout, job satisfaction	Quantitative descriptive Cross-sectional	Jordan	Ministry of Health sector of the mental health care system (one hospital and clinics).	MHNs working in the Ministry of Health sector. <i>n</i> = 130	MDS-P MBI JSS	Moral distress
Hamaideh (2011)	To examine levels of occupational stress, quality of life and	Quantitative descriptive	Jordan	All mental health care settings in Jordan.	All MHNs working in mental health settings in Jordan.	MHPSS SSS SF-36 version 2	Occupational stress

(Continued)

**TABLE 1** (Continued)

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
	social support among Jordanian mental health nurses	Correlational design			$n = 181$		
Baum and Kagan (2015)	To investigate the association between sociodemographic variables of psychiatric nurses and their intent to leave psychiatric nursing, the nursing profession and their job satisfaction, and the differences of these associations for nurses working on open and closed wards	Quantitative descriptive Cross-sectional	Israel	Open and closed psychiatric wards.	MHNs working on open and closed wards. $n = 52$	Job satisfaction questionnaire Intent to leave was measured by researcher-developed questions.	Job satisfaction Age Different inpatient settings Working part-time
Kagwe et al (2019)	To describe factors associated with job satisfaction and intent to leave among psychiatric nurses working in an inpatient state-run hospital	Quantitative descriptive Cross-sectional	United States	State-run psychiatric hospital.	All RNs working in the state-run psychiatric hospital. $n = 94$	Researcher-developed questionnaire	Salary Workload Fear of losing nursing skills Fear of assault Workplace bullying Feedback before disciplinary action Acknowledging additional credentials Opportunities for development Support Time to complete work Workplace relationships
Yanchus et al (2015)	To evaluate a model of turnover intentions across four VHA occupations: Psychologists, Psychiatrists, mental health nurses and social workers	Quantitative descriptive Cross-sectional	United States	VHA	VHA employed working in mental health services. Nurses $n = 4073$	Veterans Affairs All Employee Survey 2011	Civility Procedural justice Autonomy Psychological safety
Agyapong et al (2015)	To examine stakeholder views about the factors influencing career choices and retention of	Mixed methods Cross-sectional survey including quantitative and qualitative methods	Ghana	Three of the mental health hospitals which are integrated with mental	Psychiatrists, other doctors not specializing in psychiatry, CPNs, clinical psychologists,	Researcher-developed questionnaire.	Stigma Concerns about risk Salary

(Continued)

**TABLE 1** (Continued)

Authors and year	Aim(s)	Study design	Country	Setting	Sample and sample size	Outcome measures	Factors affecting retention identified
	community mental health workers			health outpatient facilities.	occupational therapists, social workers and other mental health workers. CPNs <i>n</i> = 71		
Gunn (2015)	To explore mental health nurses' perceptions of their career choice, and identify what factors influenced decisions to choose or remain employed in mental health nursing	Mixed methods Descriptive Cross-sectional PhD Thesis	United States	One private and one state-run hospital.	MHNs working within the two hospitals. <i>n</i> = 58	Researcher-developed questionnaire and Interviews	Confidence level Readiness to practice safely and effectively Time to spend with service users Workplace relationships Interesting and exciting nature of the job Staffing levels and ratios Salary (state facility) Burnout Returning to studies Rewarding nature of the job Stigma Personal qualities (compassion and patience)

ACS, Affective Commitment Scale; BI, Burnout Inventory; CIV, Civility; CPD, Continuing Professional Development; CPN, Community psychiatric nurse; CSE, Coping Self-efficacy scale; JSS, Job Satisfaction Survey; MBI, Maslach Burnout Inventory; MDS-P, Moral Distress Scale for Psychiatric Nurses; MHN, Mental health nurse; MHPSS, The Mental Health Professionals Stress Scale; MSQ, The Minnesota Job Satisfaction Questionnaire; NHS, National Health Service; NIOSH – The National Institute for Occupational Safety and Health; OCB, Organisational Citizenship Behaviour; POF, Person-Organisation fit scale; RN, Registered nurse; SF-36, The Short Form Health Survey; SLCI, Supervisor Leadership and Communication Inventory; SSS, The Social Support Scale; VHA, Veterans Health Administration; WBI, Work Burnout Inventory; WCS, Withdrawal Cognition Scale.

the quantitative descriptive studies met 100% of the quality criteria indicating they are high-quality studies (Hamaideh 2011; Ito *et al.* 2001; Murrels & Robinson 1999; Robinson & Tingle 2003; Robinson *et al.* 2005). Six quantitative descriptive studies included a sample that was not representative of their target population due to small sample sizes (Kagwe *et al.* 2019; Sherring & Knight 2009; Walrath 2011) and samples from single clinical settings which were not representative of MHNs working in other clinical settings in different areas (Alsaraireh *et al.* 2014; Bamber 1991; Jiang *et al.* 2019). Seven of these studies were at risk of non-

response bias judged by low response rates ranging between 35 and 65%, with no documented reasons for non-response or statistical compensation reported (Bamber 1991; Hamaideh 2014; Kagwe *et al.* 2019; Sherring & Knight 2009; Walrath 2011; Yanchus *et al.* 2015; Yanchus *et al.* 2017). Two of these studies did not use measurements that are validated or pre-tested (Bamber 1991; Kagwe *et al.* 2019). One used a measurement that was not specific to the topic under study (Yanchus *et al.* 2017). The two mixed methods studies were of low quality. Neither studies presented an adequate rationale for the use of a mixed methods design,

or any information on how or when data integration occurred. The quantitative component met 80% of the criteria for one of the studies (Agyapong *et al.* 2015), in contrast to the qualitative which met 60%, whereas in Gunn's (2015) study the qualitative component met 100% of the quality criteria and only 60% of the quantitative descriptive criteria.

## Findings

Job satisfaction was a widely applied code and cited in seven studies (Alsaraireh *et al.* 2014; Bamber 1991; Baum & Kagan 2015; Ito *et al.* 2001; Jiang *et al.* 2019; Yanchus *et al.* 2015; Yanchus *et al.* 2017). Most of these studies demonstrated that a decrease in overall job satisfaction scores was associated with an increase in intent to leave their current positions. Two of these studies demonstrated a reverse trajectory that nurses with higher overall job satisfaction scores were more likely to stay in their jobs (Bamber 1991; Jiang *et al.* 2019). Whilst it is evident that overall job satisfaction has a profound effect on intent to leave or stay, identifying the individual aspects of job satisfaction that affect retention will be more useful for directing future study and retention strategies. All identified factors were grouped into four main themes: individual characteristics, working within mental health services, training and skills and work environment.

### THEME 1: INDIVIDUAL CHARACTERISTICS

This theme is made up of three subthemes: demographic characteristics, experience and employment-related factors, and values.

#### Demographic characteristics

Age is the most significant of the demographic characteristics, with seven studies identifying an association between age and intent to leave (Alsaraireh *et al.* 2014; Bamber 1991; Baum & Kagan 2015; Ito *et al.* 2001; Jiang *et al.* 2019; Murrells & Robinson 1999; Robinson *et al.* 2005). Whilst most of the studies found that as age increases intent to leave decreases, Robinson *et al.* (2005) found that younger nurses that are aged between 20 and 29 and have children had less intentions to leave their job.

For gender, two studies found male MHNs had higher intent to leave (Alsaraireh *et al.* 2014; Jiang *et al.* 2019). Conversely, one study found no significant

difference by gender on intention to leave (Baum & Kagan 2015). Marital status was identified as a factor in two studies with MHNs with single status being more likely to intend to leave their current positions (Alsaraireh *et al.* 2014; Jiang *et al.* 2019).

Ethnicity was cited in two studies, one finding no difference between ethnicity and intent to leave (Baum & Kagan 2015) and another reporting a tentative finding that White Irish and British women were more likely to continue nursing up to ten years postqualification (Robinson *et al.* 2005). Jiang *et al.* (2019) found that MHNs with better self-rated health were less likely to leave their jobs, whilst MHNs with a higher level of education were more likely. One study found that MHNs with type A personalities were significantly more likely to leave (Bamber 1991).

These findings highlight a need to focus on how the retention of younger MHNs can be addressed and subsequently call for more research around retention and younger nurses specifically. Additionally, these findings could suggest that increasing efforts to recruit mature students into mental health nursing training may be beneficial for retention. More studies are required to form more robust conclusions on how the other demographic characteristics included in this review impact on retention.

#### Experience and employment-related factors

Bamber (1991) and Alsaraireh *et al.* (2014) found that MHNs who had left their jobs were less experienced overall than those that stayed in their jobs. This may be explained by a finding in Alexander *et al.*'s (2015) qualitative study stating:

As new [mental health nurses], participants reported starting out much more ambitious and idealistic; with experience, they had adopted a tempered realism. (Author – Alexander *et al.* 2015, p.451).

This suggests that with experience, some MHNs are able to reconsider what success means to them and their service users and feel that they are able to make positive differences to people's lives (Alexander *et al.* 2015).

The amount of time spent in their current employment was also associated with MHNs intent to leave. Two studies reported MHNs who had spent longer in their current job were more likely to stay in it (Ito *et al.* 2001; Jiang *et al.* 2019). Robinson *et al.* (2005) found MHNs who entered nursing with a degree and spent less time in their current job had greater intent

to leave, whereas those entering nursing with qualifications that were not degree sufficient and had spent less time in their current job were more likely to stay. In addition, if MHNs felt they had alternative opportunities, elsewhere they were more likely to intend on leaving their job (Ito *et al.* 2001).

The majority of new nurses are now likely to be graduates and at the beginning of their careers with limited experience as qualified registered MHNs. It is important to think about how new nurses can be better supported to spend longer in their first posts to increase their experience, so, in turn, they can support the next newly qualified nurses coming through. Future research may want to focus on the factors affecting the newly qualified nurses' intent to stay or leave their first jobs to gain more insight into their experiences, to identify any shortfalls of the preceptorship programmes that support nurses with the transition from student to qualified nurse to improve their experience and job satisfaction.

## Values

Individual values were identified in one study exploring how values affect the practice experience of MHNs (Stacey *et al.* 2011). They found that when MHNs' values conflicted with their organization's values which restricted their ability to practice in line with their own values, they contemplated leaving.

## THEME 2: WORKING WITHIN MENTAL HEALTH SERVICES

This theme is made up of three subthemes: perception of the job, factors related to practice and clinical setting.

### Perception of the job

The majority of MHNs (84%) in Gunn's (2015) study reported that the exciting and rewarding nature of the job was their reason for remaining in mental health nursing, with one MHN stating:

I have worked in many areas of nursing but this is the most rewarding job I have ever had. (Participant in Gunn 2015, p.60).

An MHN in another study also explained how the non-economic rewards of the job had contributed to their longevity in mental health nursing:

...the hook for me is seeing success, no matter how small or how large. (Participant 3 in Alexander *et al.* 2015, p.452).

Career pride was a theme that emerged through the findings of Alexander *et al.*'s (2015) study where MHNs described needing to overcome negative stereotypes held by other professionals and recognize their value, in order to develop career pride and remain in the field.

Stigma associated with working in mental health services was identified in three studies as a factor that almost always negatively affects retention of MHNs (Agyapong *et al.* 2015; Alexander *et al.* 2015; Gunn 2015). However, an MHN in Gunn's (2015) study reported that they responded to stigma by returning to the profession to care for mental health service users:

I returned to MH nursing because I am helping these patients get better than other nurses can't because they have a stigma about it'. (Participant in Gunn 2015, p.60).

### Factors related to practice

Patient-initiated violence, fear of assault and perceived risk of assault were factors identified that impact on MHNs intent to leave (Ito *et al.* 2001; Jiang *et al.* 2019; Kagwe *et al.* 2019). More than 50% of CPNs in Agyapong *et al.*'s (2015) study had contemplated leaving the profession as a result of concerns regarding risk (service user's risk to themselves or others, or risk to the MHNs themselves).

Moral distress (Hamaideh 2014; Musto & Schreiber 2012), emotional exhaustion (Sherring & Knight 2009; Yanchus *et al.* 2017) and burnout (Bamber 1991; Gunn 2015) were associated with leaving or intent to leave. MHNs reported experiencing moral distress after events that involve the safety of adolescent service users, which impacted their perceived ability to keep service users safe and caused some to leave or consider leaving their job (Musto & Schreiber 2012). One study identified that a higher level of emotional exhaustion increased the likelihood of MHNs contemplating leaving the NHS (Sherring & Knight 2009).

Being able to provide successful care and treatment, no matter how small the improvement, and remain hopeful for service users was identified in two qualitative studies (Alexander *et al.* 2015; Karłowicz & Ternus 2009). MHNs in one study describe their positive experiences of caring for service users which is what influences them to remain in mental health nursing:

'You know what keeps me there?... [Tells story of veteran with PTSD, and his family, who had a successful treatment experience]... So, to see that change, that keeps me engaged and that stays in my mind because, I think, as we go forward, there's going to be more and more and more and more people and families like his.... That's what keeps me engaged in psych. That's my story!' (Participant 4 in Alexander *et al.* 2015, p.451).

These findings suggest that working within mental health services has its own set of unique factors and factors that are more relevant to working within mental health. The rewarding nature of the job and helping people through their battles with mental health problems come at a cost to the nurses own mental and physical well-being, resulting in fear, emotional exhaustion and moral distress. It is likely that if nurses possess some of the individual factors from Theme 1, for example limited nursing experience and less time spent in their job, they will be more vulnerable to some of the factors present in this theme. This highlights the relationships between the factors across themes and the impact that one factor could have on other factors increasing the influence on MHNs' intention to leave or stay.

### Clinical setting

Interestingly, two studies reported differences in MHNs' intention to leave between different clinical settings. One study found that working on closed wards was associated with higher intent to leave mental health nursing than working on open wards in the hospital under study (Baum & Kagan 2015). Another study found that nurses working in male acute units in the hospital under study had the highest turnover intentions and MHNs working in long-term female units had the lowest turnover intentions (Alsaraireh *et al.* 2014). Unfortunately, a number of studies did not specify the individual clinical settings in which the MHNs were working, including the four studies conducted within the NHS. This means we cannot be sure which factors identified in these studies influenced retention in which settings. Due to the differences in intent to leave between clinical settings in these two studies, it would seem that factors affecting retention differ not only by organization, but also by the specific clinical settings within organizations. So, using the broad list of factors identified in this review to inform retention strategies or recommendations for change on Trust level may lead to unhelpful and unnecessary

efforts that reap few benefits, by missing what are the most important factors in which settings. With the vast range of care environments within the NHS mental health Trusts, it will be important for future studies to identify whether MHNs' intent to leave differs across clinical settings within the NHS and then explore these differences by identifying the specific factors influencing their intentions, so more tailored and meaningful retention strategies can be developed.

## THEME 3: TRAINING AND SKILLS

This theme consists of two subthemes: nurse training and developing and maintaining skills.

### Nurse training

Course preparation was a significant factor associated with plans to remain working within the NHS (Murrells & Robinson 1999). Two studies found that nurses prepared with a bachelor's degree or qualifications sufficient for degree entry were less likely to remain nursing than those with other types such as associate degrees, access courses or direct entry tests (Alsaraireh *et al.* 2014; Robinson *et al.* 2005). Beyond course preparation, one study identified that the length and quality of nurse preceptorships were factors that can cause newly qualified MHNs to consider leaving (Robinson *et al.* 2005).

### Developing and maintaining skills

Three studies demonstrated an association between MHNs' intention to leave and developing and maintaining their skills (Alexander *et al.* 2015; Kagwe *et al.* 2019; Robinson *et al.* 2005). MHNs described being part of a team that encourages and enables them to enhance their skills was a reason they remained working within the profession (Alexander *et al.* 2015; Gunn 2015). Additionally, being able to use their skills to practice autonomously was a factor that indirectly predicted MHN's turnover intentions through job satisfaction (Yanchus *et al.* 2015).

CPD and in-service training are often used by MHNs to develop and strengthen their skill sets learned during nurse training. In two studies, a lack of CPD opportunities was a contributing factor to MHN's intentions, or actual decisions, to leave their first job (Robinson *et al.* 2005; Robinson & Tingle's 2003). MHNs in Karłowicz and Ternus' (2009) study reported that they would like their organization to

promote more CPD to improve retention. In Gunn's (2015) study, CPD itself was the reason reported for why one participant left mental health nursing, as they went back to studying to further their education. Two studies identified a lack of in-service training and education negatively affects retention (Bamber 1991; Karłowicz & Ternus 2009). Whilst most of these studies found that an inability to enhance their skill set negatively affected retention, fear of losing nursing skills was also a reason cited for nurses to consider leaving their job working in a mental health hospital (Kagwe *et al.* 2019).

Whilst CPD and in-service training are important to all nurses for developing their skills, the majority of newly qualified nurses' skill development comes through working in teams with more experienced nurses who promote and encourage skill development. Although earlier findings in Theme 1 suggests that particular attention needs to be paid to the newly qualified nurses, these findings suggest that if we do not also focus on retaining experienced nurses, we may not be able to combat some of the factors affecting retention of newly qualified nurses.

#### THEME 4: WORK ENVIRONMENT

One study found that MHNs who had left their job were significantly more dissatisfied with their work environment than those still in their jobs (Bamber 1991). Work environment is the largest theme broken down into six subthemes: working relationships, leadership, organizational culture, salary, work schedule and resources.

##### Working relationships

An association between working relationships and intent to leave or stay was reported in seven studies (Alexander *et al.* 2015; Gunn 2015; Jiang *et al.* 2019; Kagwe *et al.* 2019; Karłowicz & Ternus 2009; Robinson *et al.* 2005; Robinson & Tingle 2003). Kagwe *et al.* (2019) found that all aspects of working relationships were significantly associated with the MHN's intent to leave their jobs within the next year. Karłowicz and Ternus (2009) reported mixed findings related to team dynamics with currently employed MHNs expressing positive experiences of feeling part of the team, and MHNs that had left their job reporting negative experiences and feeling outcasted. A qualitative study was able to provide a deeper understanding of what aspects of positive working relationships influence MHNs'

decisions to remain in the profession (Alexander *et al.* 2015).

Participants appreciated the sense of belonging and cohesion that the team fostered, and connected that to why they have remained in the specialty (Alexander *et al.* 2015, p.451).

Another study found that MHNs satisfied with the Trust and coordination between physicians and nurses were more likely to stay in their jobs (Jiang *et al.* 2019). In two studies, civility, defined as '*courteous and considerate workplace behaviors within the workgroup*' (Yanchus *et al.* 2015, p.222), was shown to indirectly impact on turnover intention through job satisfaction (Yanchus *et al.* 2015, 2017).

##### Leadership

Aspects of leadership were identified in seven included studies as factors that affect retention. Receiving support from line managers (Robinson *et al.* 2005), effective communication (Walrath 2011), high-quality supervision (Karłowicz & Ternus 2009) and leadership opportunities (Alexander *et al.* 2015) all aided retention.

A lack of supervisory support consisting of guidance, feedback and interpersonal support was associated with intent to leave or turnover in two studies (Ito *et al.* 2001; Yanchus *et al.* 2017). Similarly, inadequate support, especially in times of high acuity, was significantly associated with intent to leave (Kagwe *et al.* 2019). As well as line management, MHNs should have the opportunity to engage in clinical supervision for support and skill development. An unsatisfactory amount of clinical supervision was identified as a factor sufficient to cause newly qualified MHNs to contemplate leaving their job (Robinson *et al.* 2005).

##### Organizational culture

MHNs in Karłowicz and Ternus' (2009) study reported that changes within their organizations' attitude and culture should be addressed to improve retention.

Organization's procedural justice concerning performance appraisal was identified as a factor that can predict MHN's turnover intention through job satisfaction in one study (Yanchus *et al.* 2015). Likewise, being given an opportunity for feedback prior to any disciplinary action was significantly associated with lower intent to leave (Kagwe *et al.* 2019). MHNs who had left their jobs in Bamber (1991) study reported being



significantly more unsatisfied with the quality of decisions made by management.

The role of MHNs within an organization was identified as a factor that can affect retention in two studies. A role that allows MHNs to utilize their diverse set of skills (including dispensing medication) and spend a satisfactory amount of time with service users can increase job satisfaction and improve retention within the profession (Gunn 2015; Karlowicz & Ternus 2009).

Retention strategies were put in place and evaluated in two organizations committed to retaining their MHNs across two studies. Both studies found positive results with a new (HWE) orientation training programme significantly increasing retention rates for one organization (Rollins 2014); a new Nurse Residency Program for new graduates was associated with less nurse turnover for another (Pelletier *et al.* 2019).

### Salary

Salary was one of the most widely applied codes across seven studies (Agyapong *et al.* 2015; Alsaireh *et al.* 2014; Gunn 2015; Jiang *et al.* 2019; Kagwe *et al.* 2019; Karlowicz & Ternus 2009; Robinson *et al.* 2005). Jiang *et al.* (2019) found that MHNs with a higher monthly income on average are more likely to remain in their jobs. Despite job security with a permanent contract being a factor for staying in their first job, salary that does not reflect their level of responsibility was also a reason sufficient for MHNs to leave or contemplate leaving (Robinson *et al.* 2005). MHNs in Karlowicz and Ternus' (2009) study suggested pay scales and including more incentives should be addressed within their organization to improve the retention.

### Work schedule

Work schedule was explored in four studies. Shift times had no significant impact on MHN's intent to turnover (Alsaireh *et al.* 2014). MHNs working over forty hours per week (Jiang *et al.* 2019) or part-time hours (Baum & Kagan 2015) were more likely to contemplate leaving. More than a third of MHNs in Robinson *et al.*'s (2005) were dissatisfied with the combination of work hours and time spent with partners, spouses or on responsibilities for their children, and these factors were sufficient for MHNs to contemplate leaving their first mental health nursing job. MHNs in Karlowicz and Ternus' (2009) study reported that they would like scheduling and work hours to be addressed within their organization in order to improve retention.

### Resources

Newly qualified MHNs reported that unsatisfactory staffing levels, including amount of qualified to non-qualified staff, were a sufficient reason to consider leaving their first job (Robinson *et al.* 2005). Similarly, the ratio of MHNs to service users was not a reason for MHNs to remain in mental health nursing in Gunn's (2015) study.

Workload, and having adequate time to complete their work, was significantly associated with MHNs' likelihood of searching for an alternative job and intent to leave in Kagwe *et al.*'s (2019). MHNs' satisfaction with the amount of time spent on paperwork was positively related to intent to remain in the profession in Robinson *et al.* (2005).

It is clear from these findings that MHNs want to work in an organizational culture that values their role and demonstrates that through respect, procedural fairness and meaningful salary. These findings suggest that the organizational culture must also foster strong leadership, promote positive team dynamics and be sensitive to and attempt to address MHNs' concerns about the work environment such as work schedule and resources, in order to increase nurses' intent to stay in their jobs, their organization and the profession as a whole.

### DISCUSSION

This review was concerned with the individual factors that affect the retention of MHNs, and whilst overall job satisfaction is a significant factor, this concept was broken down into individual factors categorized within four interrelated themes: individual characteristics, working within mental health services, training and skills and work environment. Each of the individual factors has the potential to impact on other factors; for example, overwhelming workloads and inadequate staffing will likely expedite burnout and increase moral distress which will intensify MHNs' intention to leave. Individual factors within themes may be enough for MHNs to contemplate leaving, but due to potential intersection between thematic factors, it is likely that no factor occurs in isolation and a combination of the factors across these four themes is what contributes to overall poor job satisfaction and causes MHNs to actually leave their jobs, the NHS and the profession. The most effective retention strategies will need to address multiple yet specific factors to improve job satisfaction. MHN workforce retention will not improve on its own

or without the necessary funding to make positive changes and develop retention strategies. Therefore, policymakers and governments around the world must abandon future cuts make real investments to support the nursing profession to change the trajectory of this international crisis (Turale & Nantsupawat 2021).

Many individual and demographic characteristics are associated with MHNs' intent to leave. The most significant individual characteristic identified in this review that impacts retention is age. This finding is corroborated by the review conducted by Health Education England (2014) which found younger nurses are more likely to leave the profession. Whilst the retention of younger nurses needs to be addressed, perhaps trying to increase the uptake of mature students onto training programmes would harbour more fruitful benefits for retention. Lack of financial support is a barrier for mature students which has had a significant effect on mental health fil rates (National Health Service 2019). The findings from this review confirm the importance of NHS long-term plan's promise to explore 'earn and learn' premiums to support mature students (National Health Service 2019).

Newly qualified nurses entering their first job were found to be more at risk of leaving, which also mirrors the review by Health Education England (2014) that found turnover rates are highest in nurses' first and second year of practice. Newly qualified nurses are more likely to be younger and possess other risk factors such as less nursing experience and time spent in their current job increasing the risk of leaving for this group. Future research and retention strategies should focus on how they can better support younger and newly qualified nurses through their first two years of practice. More personalized robust preceptorships tailored to the individual nurses should be given urgent consideration.

Despite review findings suggesting older nurses are more likely to stay, one in three nurses are likely to retire over the next ten years (Royal College of Nursing 2020). Due to the global ageing workforce concern, there is a wealth of research dedicated to the retention of older nurses and preventing early retirement (Duffield *et al.* 2014; Markowski *et al.* 2020; Ryan *et al.* 2018; Uthaman *et al.* 2016). To approach this, King's College London on behalf of NHS Improvement (2019b) has set out some age-specific recommendations for the retention of older nurses and their experience which include the following: recognizing and capitalizing on their talents and ability to impart knowledge through mentoring opportunities; and promoting

flexible working (desirable shift patterns and 'wind down opportunities').

The second theme was more unique to the mental health nursing speciality. Caring for some of society's most vulnerable people with complex conditions is seen as exciting and rewarding, which is a significant factor in attracting and retaining MHNs. However, continuously caring for people with mental health problems can lead to emotional exhaustion, moral distress and burnout which negatively affect MHNs' well-being and retention. Stigma, being assaulted and perceived risk of being assaulted at work are significant factors that cause MHNs to contemplate leaving their jobs and the profession altogether. Interestingly, different clinical settings were shown to affect retention suggesting factors that affect retention differ in their existence and importance across clinical settings. Whilst there is some literature pertaining to differences in vacancy rates across different Trusts and regions, there is an absence of literature detailing retention rates for individual clinical settings within Trusts (Department of Health Social Care 2020). There is scope for future studies to explore the differences in retention rates among clinical settings within the NHS and explore what factors affecting retention exist in which clinical settings to better tailor retention strategies to provide more meaningful outcomes.

The complexities of mental health care and the vast amount of specialist services within mental health require significant training, and experience, to be able to practice competently and autonomously. Nurse training including course preparation, adequate preceptorships, sufficient in-service training and CPD opportunities is a significant factor that affects retention. Similarly, CPD was cited as a top reason for NHS staff leaving in the NHS long-term plan (National Health Service 2019). Effective training will help MHNs to hone and develop their skills and improve their confidence which will positively affect retention. It is necessary to explore in-service training, preceptorships and CPD opportunities across clinical settings to address any shortfalls. Subsequently, ascertaining what in-service training or CPD opportunities would be beneficial to MHNs in which clinical settings will better inform retention strategies.

Positive change could also be catalysed through providing preceptorship packages that are specific to the needs of the individual preceptee rather than a standardized approach. Focussing on the development of more effective preceptorships that encourage and promote CPD opportunities could be an important tool

for mitigating the risk of MHNs turning over in their first and second year of practice. Organizations need to ensure they have enough adequately experienced preceptors to meet the standard for preceptorship (Department of Health 2010), and guarantee preceptors spend enough time with their preceptee to build positive relationships and enhance person-centred skill development opportunities, which will contribute to an increase in confidence (Matua *et al.* 2014). Thereafter, regular in-service training and CPD opportunities specific to their clinical setting to enable skill development would improve MHNs' confidence and ability to work autonomously, which would enhance job satisfaction and improve retention.

Work environment constituted the largest of themes identifying multiple factors on all levels within organizations. Poor organizational cultures, limited MHN roles within the care and treatment of service users, poor leadership, low salaries in relation to responsibility, demanding work schedules, inadequate staffing, large workloads and lack of clinical supervision were factors identified that negatively impact retention. Nurses who believe they have better job opportunities elsewhere are more likely to consider leaving (Ito *et al.* 2001). This finding is in keeping with Royal College of Nursing's employment survey (Royal College of Nursing 2019) in which 55% of nurses stated they feel confident they could find a similar job with better work environments and salaries elsewhere. Work environments will vary between clinical settings, and it will be important to identify which factors are relevant in different clinical settings to understand what needs addressing to improve MHNs' work environments to improve job satisfaction and retention.

The material related to work environment alludes to a broader discussion about the nature of nursing in contemporary times. We have indicated here that poor leadership, sub-optimal supervisory structures and routines and inadequate staffing (often masked by the use of agency and bank nurses), is indicative of a system that is fragmented and open to continual change. Randall and McKeown (2014) talks of nursing being in a state of 'liquid modernity' (Bauman 2000), where care is compromised as a result of fear, uncertainty, difficulties in establishing relational continuity and the once established leadership structures now being dissipated. This review certainly indicates a workforce that is alienated and marginalized. Senek *et al.* (2020) highlight data from the adult nursing workforce where it is also clear that poor relationships with managers, an inability to solve everyday problems, transient working

relationships and perceived poor care, have resulted in alienation and intention to leave the profession.

Two retention strategies, the HWE orientation training and Nurse Residency Programme included in this review, were associated with improved retention and turnover rates of MHNs. Halter *et al.*'s (2017) review of systematic reviews on interventions to improve adult nurse turnover also illustrated the importance of some retention strategies. The National Retention Programme is creating tailored strategies on an NHS Trust level using information from exit interviews (NHS Improvement 2019a). However, MHNs turning over within the same Trust are often not considered leaving and their valuable information will not be sought or contributed to the scheme. Future studies focussed on MHNs that are still in post will elicit information that could be used to implement immediate changes or strategies that can prevent more MHNs from leaving.

## LIMITATIONS

The majority of the studies used cross-sectional designs of single hospitals at one point in time, some with small sample sizes unrepresentative of their target populations with poor response rates, limiting the generalizability of the results to other clinical settings and populations. Although a strength of this review is its inclusion of a sufficient number of studies conducted all over the world providing a global picture, different countries have different nurse training programmes and healthcare systems. This limits the generalizability of some of the findings in this review to MHNs working in the UK and more specifically the NHS.

Another limitation is that the factors identified in this review are from studies that span almost three decades. Whilst some factors from the earlier studies will still be relevant to the current workforce healthcare is an ever-changing environment in all countries which will inevitably lead to changes in the factors affecting retention of healthcare workers. Mental health nursing will have changed significantly over the past three decades in most countries. Some of the more recent changes for MHNs in the UK reported by Gournay (2005) include the following: a shift in focus within community services (community teams are caring for and treating people with more severe and enduring mental health problems than before), this has led to a shift in the patient population in inpatient services; more people in inpatient services are now more likely to be detained under the Mental Health Act and have

dual diagnoses giving rise to violence and aggression seen in these services; education and training have changed with more recent nurses likely to be graduates and more focus placed on psychosocial interventions to help equip nurses adapt to their new roles. Therefore, some of the factors in this review may not be as important for today's workforce but some may be more significant than before, for example factors relating to in-service training, CPD opportunities and risk of assault. This also highlights the need for more studies to explore the factors that affect the retention of the current workforce to develop relevant and meaningful strategies for today's workforce.

## CONCLUSION

Most themes in this review (individual characteristics, training and skills and work environment) identify similar factors that affect retention of the adult nurse workforce (Chan *et al.* 2013; Moseley *et al.* 2008). However, it is clear from this review that MHNs encounter some factors unique to working in mental health services, which suggests that retention strategies should be specific to each nursing speciality. Beyond nursing speciality, it has become apparent that the factors identified within this review are likely to vary between clinical settings in mental health due to the differences in work environments and services they provide. So, it is important that future studies now set out to explore what factors exist in which clinical settings to create more tailored retention strategies, which can then be applied on a clinical setting level across all mental health NHS Trusts, to generate better outcomes.

The majority of the studies in this review used quantitative measures that only capture the factors included in the outcome measures used, and thus, the factors identified in this review are not an exhaustive list. There is scope for studies using qualitative methods to further explore the factors affecting MHN retention to identify any important factors that may have been missed. Missing from the current literature is the voice of senior leaders who inform policy around retention. It is important to gain the perspectives of senior leaders to identify whether there are any factors affecting retention of MHNs at an organizational level that are missed by MHNs; and to explore any previous, current or planned retention strategies in order to provide tangible recommendations for change and inform effective retention strategies.

## RELEVANCE FOR CLINICAL PRACTICE

This is the first systematic review to collate all the existing evidence relating to factors that affect mental health nurse retention. The findings from this review identify and discuss some of the individual factors that affect retention and identify areas for positive change and future study. These findings can be used to inform more tailored and meaningful retention strategies for mental health NHS Trusts, with an aim to improve job satisfaction of MHNs to combat this current nursing crisis. Improving MHN retention is paramount to the safety and quality of care of our mental health service users.

## ACKNOWLEDGEMENTS

None.

## FUNDING INFORMATION

The review was supported by a grant from The Health Foundation (grant number 1275277). They provided funding for conducting the review but were not involved in the review process or decision to publish.

## REFERENCES

- Agyapong, V.I.O., Osei, A., Farren, C.K. & McAuliffe, E. (2015). Factors influencing the career choice and retention of community mental health workers in Ghana. *Human Resources for Health*, 13 (56). 10.1186/s12960-015-0050-2.
- Alexander, R.K., Diefenbeck, C.A. & Brown, C.G. (2015). Career choice and longevity in U.S. psychiatric-mental health nurses. *Issues in Mental Health Nursing*, 36 (6), 447–454.
- Alsaraireh, F., Griffin, M.T.M., Ziehm, S.R. & Fitzpatrick, J.J. (2014). Job satisfaction and turnover intention among Jordanian nurses in psychiatric units. *International Journal of Mental Health Nursing*, 23, 460–467.
- Bamber, M. (1991). Reasons for leaving among psychiatric nurses: A two-year prospective study. *Nursing Practice*, 4 (4), 9–11.
- Baum, A. & Kagan, I. (2015). Job satisfaction and intent to leave among psychiatric nurses: closed versus open wards. *Archives of Psychiatric Nursing*, 29 (4), 213–216.
- Bauman, Z. (2000). *Liquid Modernity*. Cambridge: Polity Press.
- Beech, J., Bottery, S., Charlesworth, A. *et al.* (2019). Closing the gap: Key areas for action on the health and care workforce. [Online] [Cited 28 February 2020]. Available from: URL: <https://www.kingsfund.org.uk/sites/default/file>

- s/2019-03/closing-the-gap-health-care-workforce-full-report.pdf
- British Medical Association. (2020). The impact of COVID-19 on mental health in England; Supporting services to go beyond parity of esteem. [Online] [Cited 30 September 2020]. Available from: URL: <https://www.bma.org.uk/media/2750/bma-the-impact-of-covid-19-on-mental-health-in-england.pdf>
- Buchan, J., Charlesworth, A., Gershlick, B. & Seccombe, I. (2019). A critical moment: NHS staffing trends, retention and attrition. [Online] [Cited 28 February 2020]. Available from: URL: [https://www.health.org.uk/sites/default/files/upload/publications/2019/A%20Critical%20Moment\\_1.pdf](https://www.health.org.uk/sites/default/files/upload/publications/2019/A%20Critical%20Moment_1.pdf)
- Buchan, J., Twigg, D., Dussault, G., Duffield, C. & Stone, P.W. (2015). Policies to sustain the nursing workforce: an international perspective. *International Nursing Review*, 62, 162–170.
- Care Quality Commission. (2017). The state of care in mental health services 2014 to 2017: Findings from CQC's programme of comprehensive inspections of specialist mental health services. [Online] [Cited 24 March 2020]. Available from: URL: [https://www.cqc.org.uk/sites/default/files/20170720\\_stateofmh\\_report.pdf](https://www.cqc.org.uk/sites/default/files/20170720_stateofmh_report.pdf)
- Chan, Z.C.Y., Tam, W.S., Lung, M.K.Y., Wong, W.Y. & Chau, C.W. (2013). A systematic literature review of nurse shortage and the intention to leave. *Journal of Nursing Management*, 12, 605–613.
- Department of Health (2010). Preceptorship Framework: for Newly Registered Nurses, Midwives and Allied Health Professionals. [Online] [Cited 23 December 2020]. Available from: URL: [https://webarchive.nationalarchives.gov.uk/20130105024308/http://www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/@dh/@en/@about/documents/digitalasset/dh\\_1141116.pdf](https://webarchive.nationalarchives.gov.uk/20130105024308/http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/@about/documents/digitalasset/dh_1141116.pdf)
- Department of Health and Social Care. (2020) The NHS nursing workforce. [Online] [Cited 23 December 2020]. Available from: URL: <https://www.nao.org.uk/wp-content/uploads/2020/03/The-NHS-nursing-workforce-Summary.pdf>
- Dewanto, A. & Wardhani, V. (2018). Nurse turnover and perceived causes and consequences: a preliminary study at private hospitals in Indonesia. *BMC Nursing*, 17 (2), 52.
- Dixon-Woods, M., Agarwal, S., Jones, D., Young, B. & Sutton, A. (2005). Synthesising qualitative and quantitative evidence: A review of possible methods. *Journal of Health Services Research & Policy*, 10 (1), 45–53.
- Duffield, C., Graham, E., Donoghue, J., Griffiths, R., Bichel-Findlay, J. & Dimitrelis, S. (2014). Why older nurses leave the workforce and the implications of them staying. *Journal of Clinical Nursing*, 24, 824–831.
- Gournay, K. (2005). The changing face of psychiatric nursing: Revisiting mental health nursing. *Advances in Psychiatric Treatment*, 11, 6–11.
- Gunn, R. (2015). *Mental Health Nurses' Perceptions: Making a Career Choice and Remaining in Mental Health Nursing*. ProQuest, LLC:10605386.
- Halter, M., Pelone, F., Boiko, O. et al. (2017). Interventions to Reduce Adult Nursing Turnover: A Systematic Review of Systematic Reviews. *Open Nursing Journal*, 11, 108–123.
- Hamaideh, S.H. (2011). Occupational Stress, Social Support, and Quality of Life among Jordanian Mental Health Nurses. *Issues in Mental Health Nursing*, 33 (1), 15–23.
- Hamaideh, S.H. (2014). Moral distress and its correlates among mental health nurses in Jordan. *International Journal of Mental Health Nursing*, 23, 33–41.
- Hanohano, C. & Dee, V. (2017). Physical assault, perceived stress, coping and attitudes toward assault experienced by psychiatric nurses and their intent to leave. *Madridge Journal of Nursing*. <https://doi.org/10.18689/2638-1605.a2.002>.
- Happell, B. (2008). Exploring workforce issues in mental health nursing. *Contemporary Nurse*, 29 (1), 43–51.
- Health Education England. (2014). Growing Nursing Numbers: Literature review on nurses leaving the NHS. [Online] [Cited 29 March 2020]. Available from: URL: <https://www.hee.nhs.uk/sites/default/files/documents/Nurses%20leaving%20practice%20-%20Literature%20Review.pdf>
- Heyvaert, M., Hannes, K. & Onghena, P. (2016). *Using Mixed Methods Research Synthesis for Literature Reviews*. Los Angeles: SAGE.
- Hong, Q.N., Pluye, P., Fabregues, S. et al. (2018). Mixed Methods Appraisal Tool (MMAT) Version 2018: User guide. [Online] [Cited 02 November 2020]. Available from: URL: [http://mixedmethodsappraisaltoolpublic.pbworks.com/w/file/attach/127916259/MMAT\\_2018\\_criteria-manual\\_2018-08-01\\_ENG.pdf](http://mixedmethodsappraisaltoolpublic.pbworks.com/w/file/attach/127916259/MMAT_2018_criteria-manual_2018-08-01_ENG.pdf)
- Ito, H., Eisen, S.V., Sederer, L.I., Yamada, O. & Tachimori, H. (2001). Factors affecting psychiatric nurses' intention to leave their current job. *Psychiatric Services*, 52 (2), 232–234.
- Jiang, F., Zhou, H., Rakofsky, J. et al. (2019). Intention to leave and associated factors among psychiatric nurses in China: A nationwide cross-sectional study. *International Journal of Nursing Studies*, 94, 159–165.
- Jones, C.B. & Gates, M. (2007). The Cost and Benefits of Nurse Turnover: A Business Case for Nurse Retention. *The Online Journal of Issues in Nursing*, 12 (3), <https://doi.org/10.3912/OJIN.Vol12No03Man04>.
- Kagwe, J., Jones, S. & Johnson, S.L. (2019). Factors related to intention to leave and job satisfaction among registered nurses at a large psychiatric hospital. *Issues in Mental Health Nursing*, 40 (9), 754–759.
- Karłowicz, K.A. & Ternus, M.P. (2009). Issues influencing psychiatric nurse retention during the first year of employment: a case analysis. *Journal of Nursing Management*, 17, 49–58.
- Markowski, M., Cleaver, K. & Weldon, S.M. (2020). An integrative review of the factors influencing older nurses' timing of retirement. *Journal of Advanced Nursing*, 76 (9). <https://doi.org/10.1111/jan.14442>.

- Matua, G.A., Seshan, V., Savithri, R. & Fronda, D.C. (2014). Challenges and strategies for building and maintaining effective preceptor-preceptee relationships among nurses. *Sultan Qaboos University Medical Journal*, 14 (4), e530–536.
- Moher, D., Liberati, A., Tetzlaff, J. & Altman, D.G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA Statement. *Open Medicine*, 3 (3), 123–130.
- Moseley, A., Jeffers, L. & Paterson, J. (2008). The retention of the older nursing workforce: A literature review exploring factors that influence the retention and turnover of older nurses. *Contemporary Nurse*, 30 (1), 46–56.
- Murrells, T. & Robinson, S. (1999). Phase 1 of a longitudinal study of mental health nurses' careers: Exploring relationships between profiles, course perceptions and plans to work in the NHS. *Journal of Mental Health*, 8 (1), 55–68.
- Musto, L. & Schreiber, R.S. (2012). Doing the best i can do: Moral distress in adolescent mental health nursing. *Issues in Mental Health Nursing*, 33 (3), 137–144.
- National Health Service. (2019). Online version of the NHS Long Term Plan – Chapter 4: NHS staff will get the backing they need – 5. Supporting our current NHS staff. [Online] [Cited 13 July 2020]. Available from: URL: <https://www.longtermplan.nhs.uk/online-version/chapter-4-nhs-staff-will-get-the-backing-they-need/5-supporting-our-current-nhs-staff/>
- NHS Improvement. (2019a). The national retention programme: two years on. [Online] [Cited 31 March 2020]. Available from: URL: <https://improvement.nhs.uk/resources/national-retention-programme-two-years-on/#content>
- NHS Improvement (2019b) Retention of older nurses: a focus group study in UK hospitals. [Online]. [Cited 17 December 2020]. Available from: URL: [https://improvement.nhs.uk/documents/4938/Retention\\_of\\_older\\_nurses\\_-\\_exec\\_summary\\_.pdf](https://improvement.nhs.uk/documents/4938/Retention_of_older_nurses_-_exec_summary_.pdf)
- Nursing & Midwifery Council (2017). The NMC register. [Online] [Cited 28 February 2020]. Available from: URL: <https://www.nmc.org.uk/globalassets/sitedocuments/other-publications/nmc-register-2013-2017.pdf>
- O'Brien-Pallas, L., Murphy, G.T., Shamian, J., Li, X. & Hayes, L.J. (2010). Impact of determinants of nurse turnover: a pan-Canadian study. *Journal of Nursing Management*, 18, 1073–1086.
- Pelletier, L.R., Vincent, C., Woods, L., Odell, C. & Stichler, J.F. (2019). Effectiveness of a psychiatric-mental health nurse residency program on retention. *Journal of the American Psychiatric Nurses Association*, 25 (1), 66–75.
- Pluye, P. & Hong, Q.N. (2014). Combining the power of stories and the power of numbers: Mixed methods research and mixed studies reviews. *The Annual Review of Public Health*, 35, 29–45.
- Randall, D. & McKeown, M. (2014). Failure to care: nursing in a state of liquid modernity? *Journal of Clinical Nursing*, 23 (5–6), 766–767.
- Robinson, S., Murrells, T. & Smith, E.M. (2005). Retaining the mental health nursing workforce: Early indicators of retention and attrition. *International Journal of Mental Health Nursing*, 14, 230–242.
- Robinson, S. & Tingle, A. (2003). Continuing education opportunities for recently qualified mental health diplomates. *Journal of Psychiatric and Mental Health Nursing*, 10 (6), 659–668.
- Rollins, L. (2014). Healthy Work Environment Orientation Training and Psychiatric Nurse Retention. Walden Dissertations and Doctoral Studies Collection. [Cited 08 June 2020]. Available from: URL: [https://pdfs.semanticscholar.org/f4ad/5afb3d4e675b5a6d3610f55c30048ed6f1dc.pdf?\\_ga=2.69834448.1315673077.1588336503-975706568.1587993464](https://pdfs.semanticscholar.org/f4ad/5afb3d4e675b5a6d3610f55c30048ed6f1dc.pdf?_ga=2.69834448.1315673077.1588336503-975706568.1587993464)
- Royal College of Nursing (2015). A workforce in crisis? The UK nursing labour market review 2015. [Online] [Cited 24 March 2020]. Available from: URL: <https://www.rcn.org.uk/professionaldevelopment/publications/pub-005348>
- Royal College of Nursing. (2019). RCN Employment Survey 2019. [Online] [Cited 08 April 2020]. Available from: URL: <https://www.rcn.org.uk/professional-development/publications/pub-007927>
- Royal College of Nursing. (2020) Nursing Shortages in England: Westminster Hall Debate, Tuesday 3 March, 9.30–11am. [Online] [Cited 18 December 2020]. Available from: URL: [https://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwi9z\\_P1xtftAhVxmVwKHRiBAkAQFjAAegQIAxAC&url=https%3A%2F%2Fwww.rcn.org.uk%2F-%2Fmedia%2Froyal-college-of-nursing%2Fdocuments%2Fpublications%2F2020%2Fmarch%2F009-174.pdf%3F%3Den&usq=AOvVaw0WjzLVOLRs9mgP6zGC3vJx](https://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKEwi9z_P1xtftAhVxmVwKHRiBAkAQFjAAegQIAxAC&url=https%3A%2F%2Fwww.rcn.org.uk%2F-%2Fmedia%2Froyal-college-of-nursing%2Fdocuments%2Fpublications%2F2020%2Fmarch%2F009-174.pdf%3F%3Den&usq=AOvVaw0WjzLVOLRs9mgP6zGC3vJx)
- Ryan, C., Bergin, M., White, M. & Wells, J.S.G. (2018). Ageing in the nursing workforce – a global challenge in an Irish context. *International Nursing Review*, 66 (2). <https://doi.org/10.1111/inr.12482>.
- Senek, M., Robertson, S., Ryan, T. *et al.* (2020). Determinants of nurse job dissatisfaction-findings from a cross-sectional survey analysis in the UK. *BMC Nursing*, 19 (1), 1–10.
- Sherring, S. & Knight, D. (2009). An exploration of burnout among city mental health nurses. *British Journal of Nursing*, 18 (20), 1234–1240.
- Stacey, G., Johnston, K., Stickley, T. & Diamond, B. (2011). How do nurses cope when values and practice conflict? *Nursing Times*, 107 (5), 20–23.
- Thomas, J. & Harden, A. (2008). Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Medical Research Methodology*, 8 (45). <https://doi.org/10.1186/1471-2288-8-45>.
- Turale, S. & Nantsupawat, A. (2021). Clinician mental health, nursing shortage and the COVID-19 pandemic: Crises within crises. *International Nursing Review*, 68 (1), 12–14.
- Unison. (2017a). Struggling to Cope: Mental Health Staff and Services Under Pressure. [Online] [Cited 08 July 2020].

- Available from: URL: <https://www.unison.org.uk/content/uploads/2017/10/Struggling-to-Cope.pdf>
- Unison. (2017b). Ratios not Rationing: Unison safe staffing report 2017. [Online] [Cited 24 March 2020]. Available from: URL: <https://www.unison.org.uk/content/uploads/2017/04/Rationotrationing.pdf>
- Uthaman, T., Chua, T.L. & Ang, S.Y. (2016). Older nurses: A literature review on challenges, factors in early retirement and workforce retention. *Proceedings of Singapore Healthcare*, 25 (1), 50–55.
- Walrath, D. (2011). *A Quantitative Correlational Study: Nurse Leadership and Communication Effect on Psychiatric Nurse Performance and Retention*. Unpublished PhD thesis. University of Phoenix.
- Ward, L. (2011). Mental health nursing and stress: Maintaining balance. *International Journal of Mental Health Nursing*, 20 (2), 75–85.
- World Health Organization (2013). A universal truth: No health without a workforce. [Online] [Cited 28 February 2020]. Available from: URL: [https://www.who.int/workforcealliance/knowledge/resources/GHWA-a\\_universal\\_truth\\_report.pdf?ua=1](https://www.who.int/workforcealliance/knowledge/resources/GHWA-a_universal_truth_report.pdf?ua=1)
- World Health Organization. (2020). Nursing and Midwifery: Key facts. [Online] [Cited 19 May 2020]. Available from: URL: <https://www.who.int/news-room/fact-sheets/detail/nursing-and-midwifery>.
- Yanchus, N.J., Periard, D., Moore, S.C., Carle, A.C. & Osatuke, K. (2015). Predictors of job satisfaction and turnover intention in VHA mental health employees: A comparison between psychiatrists, psychologists, social workers, and mental health nurses. *Human Service Organizations: Management, Leadership & Governance*, 39 (3), 219–244.
- Yanchus, N.J., Periard, D. & Osatuke, K. (2017). Further examination of predictors of turnover intention among mental health professionals. *Journal of Psychiatric and Mental Health Nursing*, 24, 41–56.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Supplementary Material